

Trait-based diet selection: prey behaviour and morphology predict vulnerability to predation in reef fish communities

Stephanie J. Green^{1,2*} and Isabelle M. Côté¹

¹Department of Biological Sciences, Simon Fraser University, 8888 University Drive, Burnaby, BC V5A 1S6, Canada; and ²Department of Zoology, Oregon State University, Corvallis, OR 97331-2914, USA

Summary

1. Understanding how predators select their prey can provide important insights into community structure and dynamics. However, the suite of prey species available to a predator is often spatially and temporally variable. As a result, species-specific selectivity data are of limited use for predicting novel predator–prey interactions because they are assemblage specific.

2. We present a method for predicting diet selection that is applicable across prey assemblages, based on identifying general morphological and behavioural traits of prey that confer vulnerability to predation independent of species identity. We apply this trait-based approach to examining prey selection by Indo-Pacific lionfish (*Pterois volitans* and *Pterois miles*), invasive predators that prey upon species-rich reef fish communities and are rapidly spreading across the western Atlantic.

3. We first generate hypotheses about morphological and behavioural traits recurring across fish species that could facilitate or deter predation by lionfish. Constructing generalized linear mixed-effects models that account for relatedness among prey taxa, we test whether these traits predict patterns of diet selection by lionfish within two independent data sets collected at different spatial scales: (i) *in situ* visual observations of prey consumption and availability for individual lionfish and (ii) comparisons of prey abundance in lionfish stomach contents to availability on invaded reefs at large.

4. Both analyses reveal that a number of traits predicted to affect vulnerability to predation, including body size, body shape, position in the water column and aggregation behaviour, are important determinants of diet selection by lionfish. Small, shallow-bodied, solitary fishes found resting on or just above reefs are the most vulnerable. Fishes that exhibit parasite cleaning behaviour experience a significantly lower risk of predation than non-cleaning fishes, and fishes that are nocturnally active are at significantly greater risk. Together, vulnerable traits heighten the risk of predation by a factor of nearly 200.

5. Our study reveals that a trait-based approach yields insights into predator–prey interactions that are robust across prey assemblages. Importantly, *in situ* observations of selection yield similar results to broadscale comparisons of prey use and availability, which are more typically gathered for predator species. A trait-based approach could therefore be of use across predator species and ecosystems to predict the outcomes of changing predator–prey interactions on community dynamics.

Key-words: foraging behaviour, *in situ* observations, predator–prey interactions, prey characteristics, *Pterois volitans/miles*, resource use and availability, selective predation, stomach contents analysis

*Correspondence author. E-mail: steph.j.green@gmail.com

Introduction

Predation affects the demographic rates of species in most ecosystems (Caley 1993; Carr & Hixon 1995; Arcese, Smith & Hatch 1996; Lima *et al.* 2001; Creel *et al.* 2007). Understanding the process by which predators select their prey can therefore provide key insights into the structure and function of communities (Gilinsky 1984; Holt 1984; Kotler 1984; Karanth & Sunquist 1995; Juanes, Buckel & Scharf 2001; Almany & Webster 2004). Optimal foraging theory predicts that predators should select prey items in a manner that maximizes energetic gain while minimizing the energy expended in locating, handling and processing the item (Pyke, Pulliam & Charnov 1977). If all prey types are equally profitable, then predators should consume prey in proportion to their abundance in the environment, so diet composition will track changes in prey community composition across space and time (e.g. Beukers-Stewart & Jones 2004). However, predators often consume some prey in quantities disproportionate to their availability (i.e. dietary preference or selective predation; e.g. Green & Osborne 1981; Scott & Murdoch 1983; Donázar & Ceballos 1989; Hambright 1991; Hernández 1995; Almany *et al.* 2007; Nunn, Tweson & Cowx 2012).

Identifying prey species that are selectively consumed (or, conversely, avoided) has been the aim of much foraging ecology research, with a large number of studies focused on which species predators choose in experimental settings (e.g. Einfalt & Wahl 1997; Nilsson & Brönmark 2000; Scharf *et al.* 2003; Heinonen & Auster 2012; Provost *et al.* 2006), and a smaller number focused on species selection *in situ* (e.g. Cunningham, Gustavson & Ballard 1999; Isaac *et al.* 2012; Wen *et al.* 2012). However, the assemblage of prey available to a given predator is spatially and temporally variable, owing to both natural processes [e.g. gradients in prey assemblage across the predator's range (e.g. Holt 1984; Steele 1989; Bost *et al.* 2002; Phillips *et al.* 2009; Jaeger *et al.* 2010)] and, increasingly, anthropogenically mediated species additions [i.e. through invasion; e.g. (Caut, Angulo & Courchamp 2008)] and removals [e.g. through exploitation; e.g. (Crooks & Soulé 1999; Myers & Worm 2003)], limiting the usefulness of experimental and assemblage-specific findings for predicting selection from varying sets of prey.

The heterogeneous nature of predator–prey interactions necessitates an approach to predicting diet selection that can be applied across changing community compositions. Selective predation is thought to occur when prey types vary in one or more morphological or behavioural characteristics that affect their encounter rate and handling time (Pyke, Pulliam & Charnov 1977). It therefore follows that diet selection should be predictable across prey assemblages if vulnerability to predation is conferred by morphological and/or behavioural traits of prey that recur across species assemblages. Thus, the strength of interactions when a predator is exposed to new prey types could be predicted by identifying traits associated with vulnera-

bility to predation, independently from species identity. A functional, or trait-based, approach has been advocated as a means to make community ecology more predictive (McGill *et al.* 2006), but we are not aware of studies that have tested this idea in the context of predator–prey interactions.

To illustrate a trait-based approach to studying diet selection, we evaluate whether morphological and behavioural characteristics of Caribbean coral reef fishes affect vulnerability to predation by the Indo-Pacific lionfish (*Pterois volitans* and *Pterois miles*). Lionfish have rapidly become established on coral reefs across the western Atlantic, Caribbean, and Gulf of Mexico, and the invasion continues to spread southwards (Betancur *et al.* 2011; Côté, Green & Hixon 2013b; Côté *et al.* 2013a). Predation by invasive lionfish has been implicated in the significant decline of reef fish biomass in the northern Caribbean (Albins and Hixon 2008; Green *et al.* 2012, in press). Given that reef fish assemblages differ across this large region, elucidating prey trait variants that are most susceptible to predation will aid in forecasting prey declines as the invasion spreads to areas with novel fish assemblages. To do this, we generated hypotheses about prey traits that could facilitate or deter predation by lionfish, and assessed whether selective predation on specific trait variants occurs from data collected at two spatial scales. At the smallest scale, we simultaneously recorded prey available to and prey consumed by individual lionfish hunting. At a larger scale, we compared the abundance of prey items in lionfish stomach contents to prey abundance on invaded reefs at large during underwater visual observations.

Materials and methods

PREY TRAIT HYPOTHESES

To simultaneously examine the effects of prey morphology and behaviour on lionfish diet selection, we compiled information on a suite of traits that are likely to influence vulnerability to predation (Table 1) and generated hypotheses for each. We then assigned a score or value for each trait to each fish species encountered on our study reefs (Table S1, Supporting information), based on published descriptions of each species (Böhlke & Chaplin 1993; Deloach 1999; FAO 2002a,b; Humann & Deloach 2002).

Behaviour

Lionfish are visual predators that employ a stalking strategy, in which they approach their prey by hovering slowly over the substrate (Côté & Maljković 2010; Green, Akins & Côté 2011). Several aspects of prey behaviour should confer variable vulnerability to this hunting style. We predicted that demersal (i.e. living <2 m from the bottom) fishes would be more vulnerable to predation than benthic species (i.e. those living on or in the benthos), which may be less visible, and that pelagic fishes occupying the water column above reefs (i.e. >2 m above the

Table 1. Morphological and behavioural traits of prey fish predicted to influence vulnerability to predation by invasive lionfish (*Pterois volitans/miles*)

	Prey attribute	Data type	Units/levels	Predictions of vulnerability to predation
Morphology	Body shape	Continuous	Body length to depth ratio: Higher values indicate increasingly linear shapes	↑ ratio (i.e. ↑ shallow bodied) = ↑ vulnerability
	Body size	Continuous	Mass (g)	↑ mass = ↓ vulnerability
	Physical or chemical defence	Binary	Yes/no = 0/1	+ defence = ↓ vulnerability
Behaviour	Cleaning behaviour	Binary	Yes/no = 0/1	+ cleaning behaviour = ↓ vulnerability
	Aggregation size	Categorical	Solitary, shoaling, schooling	Solitary < shoaling < schooling
	Nocturnally active	Binary	Yes/no = 0/1	+ nocturnality = ↑ vulnerability
	Water column position	Categorical	Benthic, demersal, pelagic	Pelagic < benthic < demersal

benthos) would be least accessible. While many studies suggest that schooling behaviour is an effective anti-predator strategy because aggregations can dilute individual risk of predation (Pitcher & Parrish 1993), we hypothesized that lionfish, whose slow movements afford them a close approach to potential prey, may in fact target schooling fish species because the conspicuousness of schools may reduce predator search time. We recognized three categories of aggregation behaviour: solitary, shoaling and schooling. We defined shoaling as an intermediate state, based on the observation that many fish species tend to form loose feeding aggregations as juveniles (e.g. wrasses and parrotfishes). We also hypothesized that fishes that are nocturnally active and whose emergence from sheltered reef habitats overlaps with the timing of lionfish foraging behaviour at crepuscular times (Green, Akins & Côté 2011) would be more vulnerable to lionfish predation than their diurnal counterparts because this strategy may result in increased encounters with hunting lionfish (Hobson 1973; Fishelson 1975; Green, Akins & Côté 2011). Finally, we anticipated that fishes that exhibit cleaning behaviour, either facultatively (i.e. only as juvenile) or obligately (i.e. throughout their lifespan), may be less readily consumed by lionfish, because such species often experience lower mortality rates from predation, possibly because of recognition by fish predators of the service cleaners provide (e.g. Colin 1975; Côté 2000).

Morphology

We considered three major aspects of prey morphology that may affect vulnerability to predation by lionfish. For gape-limited predators, the costs of prey consumption (in terms of energy and time for capture and handling) increase with prey body depth (Hambricht 1991; Nilsson & Brönmark 2000). We thus anticipated that prey body shape plays an important role in prey selection by gape-limited lionfish, with vulnerability to predation decreasing with increasingly deeper-bodied (i.e. increasingly round) shapes. We quantified shape as the ratio of total length (TL) to maximum body depth, such that higher values indicate less round (i.e. more elongate), and hence more vulnerable, shapes. We calculated the average ratio of fish TL to body depth (in cm), measured as the linear distance from the pelvic girdle to the dorsal ridge, as in Hambricht (1991), from lateral images of 3–5 specimens [available in Humann & Deloach (2002) and on FishBase (www.fishbase.org)] of each fish species encountered on the study reefs. Images were only used for specimens that were within a size range that could be consumed by lionfish (i.e. <15 cm TL; Green *et al.* 2012). Second, we hypothesized that

fishes possessing a chemical or physical defence (e.g. a toxic or distasteful compound, sharp spine(s) or barb) would be less vulnerable to predation compared with their undefended counterparts (e.g. Hoogland, Morris & Tinbergen 1956; Caley & Schluter 2003; Schubert *et al.* 2003). Finally, the average size of prey (TL) relative to the size of their lionfish predator across several studies of lionfish stomach contents is ~15%; much smaller than the maximum of ~48% (Morris & Akins 2009; Munoz, Currin & Whitfield 2011; Côté, Green & Hixon 2013b; Côté *et al.* 2013a). It is unclear whether this result simply reflects the abundance of various prey sizes on invaded reefs. However, there is ample experimental evidence that gape-limited predators tend to selectively consume prey that are far smaller than their maximum gape limits, likely because larger prey are better at evading predators and smaller prey are less costly to capture and handle by predators (Einfalt & Wahl 1997; Nilsson & Brönmark 2000). We therefore hypothesized that vulnerability to predation would decrease with increasing prey size, quantified as body mass (in g), which we obtained by converting estimates of TL (to the nearest 1 cm) to weight using the allometric scaling function mass (g) = $a \cdot TL(\text{cm})^b$, where a and b are species-specific constants that relate length to weight obtained from FishBase (www.fishbase.org).

IN SITU OBSERVATIONS OF PREY SELECTION

To quantify lionfish prey selection *in situ*, we conducted detailed visual observations of lionfish on shallow coral patch reefs off Eleuthera Island, the Bahamas (24° 50.000 N, 76° 14.000 W), in September 2008 and December 2009. On invaded reefs, lionfish exhibit bold behaviour and have little fear of divers, which facilitates a close approach without altering behaviour (Côté & Maljković 2010; Green, Akins & Côté 2011). They hunt prey via a slow, hovering, stalking strategy, which affords observers easy view of the target prey, as well as the success of strikes at prey (Côté & Maljković 2010; Green, Akins & Côté 2011). Individual lionfish were observed continuously for 30 min by a scuba diver from a distance of 2–3 m (as per Green, Akins & Côté 2011). Only individuals that were actively hunting (i.e. oriented with head down, pectoral fins spread and actively stalking prey fishes) and consumed at least one prey item during the 30-min observation period were included in this study. At the onset of observations, we estimated the size of the focal lionfish (TL to the nearest 1 cm). During each observation period, we recorded the identity, abundance and size (TL to the nearest 1 cm) of all fishes within a 1-m radius of the hunting lionfish

every 5 min (hence, six point counts per observation). The large number of fish precluded continuous tabulation of prey across the 30-min period. We also recorded the identity and size of all prey items consumed by lionfish during the 30-min observation period, as well as the fishes within a 1-m radius of the focal lionfish immediately following each prey capture. Lionfish are visual predators that primarily hunt in the open over the top of coral reefs or sand/sea grass beds. We thus assumed that only prey within visual sight of the lionfish (i.e. not those in crevices or under ledges) were available for consumption. We classified all fishes of <15 cm TL as potential prey items, based on physiological limits on predator: prey size ratio set by gape size (Nilsson & Brönmark 2000).

PREY SELECTION FROM STOMACH CONTENTS ANALYSIS

We also inferred lionfish prey selection by comparing the abundance of prey items in the stomachs of lionfish collected from eight sites along a continuous coral reef system off southwest New Providence Island, the Bahamas (24°59.072 N, 77°32.207 W), to the abundance of prey identified during visual surveys of the same sites. Lionfish collections and prey visual surveys were conducted at depths of 10–20 m between May 2008 and July 2008. Lionfish were collected using hand nets and euthanized in a clove oil and seawater solution at the surface. We then measured TL to the nearest 1 cm, extracted stomach contents from each specimen, and identified all fish prey items visually to the lowest taxonomic resolutions possible. For whole fish prey (i.e. items for which TL could be estimated) that could not be identified to species because of degradation of key skin pigments and coloration, we obtained species identity via DNA barcode analysis (Côté *et al.* 2013a). To minimize potential identification bias, we only included in our analyses lionfish stomachs from which all of the fish items could be identified to species (either visually or via DNA barcoding).

To estimate the abundance of prey available to lionfish, we conducted detailed surveys of prey-sized fishes (i.e. <15 cm TL) on 6–12 belt transects (30 m × 2 m) at each of the eight lionfish capture locations. Along each transect, a trained observer carefully searched in all holes and crevices for cryptic fishes, using a dive light as needed. The identity and TL of all fishes were recorded to the nearest 1 cm. We assumed that any prey hidden within the reef framework and not recorded in our visual surveys would not be available to lionfish to consume.

Transect surveys of prey-sized fishes were conducted immediately prior to lionfish collections at each site. We assumed that the assemblage of prey-sized fishes observed on our transect surveys matched the assemblage of prey fishes available to lionfish during hunting bouts at the sites within ~24 h of capture, based on laboratory-derived times to digestion for lionfish prey (J.A. Morris, unpublished data) and preliminary results from external tagging and tracking of lionfish that suggested limited movement on the study reefs (S.J. Green, unpublished data).

STATISTICAL ANALYSIS

In the absence of prey selection, our null expectation was that lionfish should consume prey types in proportion to their abundance. Our analyses of direct and indirect observations of lionfish predation tested whether the consumption by lionfish of prey

types varying in morphology and behaviour deviated significantly from this null expectation. Specifically, we used generalized linear mixed-effects models (GLMMs) to assess the effects of prey morphology and behaviour on lionfish prey selection from our direct and indirect observations of predation. We created models for each data set (i.e. direct and indirect observations) using the `glmer()` function in the package `lme4` for the statistical software `R` (Bates, Maechler & Bolker 2012; R Core Team 2012). Because morphological and behavioural traits can be shared among related species and are thus not statistically independent (e.g. Jennings *et al.* 1999), we accounted for phylogenetic relatedness among prey fishes by specifying family as a nested random effect in our models. In the absence of a complete molecular phylogeny for the reef fishes in our study region, we classified species based on their most recent taxonomic groupings obtained from Fish-Base (www.fishbase.org), cross-checked with the World Registry of Marine Species (www.marinespecies.org).

In situ observations

To identify the drivers of lionfish prey selection from our direct observations of lionfish hunting on Eleuthera reefs, we first quantified the abundance of potential prey types [species and TL (in 1-cm bins) combination] within 1 m of focal hunting lionfish and then calculated the proportion of each type that was consumed (i.e. proportional response variable). As potential predictors of prey selection (proportion consumed), we included individual prey size (mass in g), species-specific body shape (quantified as the ratio of body length to depth), position in the water column, whether the prey species was nocturnal, physically or chemically defended, and a cleaner (fixed effects; Table 1). In addition to fish family, we specified three nested random effects, so that each potential prey was nested within the point count on which it was observed, the focal lionfish that was being observed, and the body length of the lionfish (TL to the nearest 1 cm). Parameter estimates for each fixed and random effect were obtained using the Laplace approximation with a binomially distributed error and cloglog-link function because the proportion of zeros in our data set greatly exceeded the proportion of ones (Bolker *et al.* 2009; Zuur *et al.* 2009). We conducted multiple Bonferroni-corrected Wald Z tests to obtain pairwise comparisons of the relative selection probabilities for all levels of all categorical variables.

Stomach contents analysis

To identify the drivers of lionfish prey selection from our indirect observations of lionfish stomach contents and prey availability on New Providence reefs, we first calculated the proportion of each fish prey type (species and length class combinations, in 1-cm bins) consumed by lionfish by first summing the abundance of each type across the stomach contents of all lionfish, and then dividing by its abundance estimate from our visual surveys of availability on the reefs. Next, we constructed a GLMM where the proportion of each prey type consumed (i.e. proportional response) was predicted by its morphological and behavioural characteristics (fixed effects; Table 1) and weighted by its abundance in surveys of the study area (following Zuur *et al.* 2009). Again, we conducted multiple Bonferroni-corrected post hoc Wald Z tests to obtain pairwise comparisons of the relative selection probabilities for each categorical predictor.

Model selection and multi-model averaging

For both of our analyses (direct and indirect observations), we evaluated the relative strength of support for predictions about the effect of prey morphological and behavioural traits, and incorporated model uncertainty using AICc (for small sample sizes) model selection and averaging, respectively (Burnham & Anderson 2002). AICc values represent the trade-off between model complexity (i.e. the number of parameters) and model fit within a candidate set of models. In order to compare relative effect sizes, we standardized our data by centring (subtracting the mean) and dividing by two standard deviations (Gelman 2008). We calculated average parameter estimates for each prey trait variable from its weights in the set of top models (i.e. those with $\Delta\text{AICc} < 4$) in the MuMIn package (Barton 2013) for R statistical software, using the 'Natural Average' method (Burnham & Anderson 2002; Grueber *et al.* 2011).

Results

PREY SELECTION FROM *IN SITU* OBSERVATIONS

On Eleuthera reefs, we observed 22 hunting lionfish capture a total of 32 prey fishes from 12 species and six families (Table S1, Supporting information). Lionfish size ranged from 10 to 36 cm TL (26 ± 6 cm; mean \pm SD), while their captured prey ranged from 2 to 7 cm TL (4 ± 1 cm; mean \pm SD). The maximum number of prey consumed by a single lionfish during our observations was 4; 12 lionfish consumed a single prey fish. There were 32 species from 16 families recorded in point counts of fish prey availability (i.e. all individuals within a 1-m radius of the focal lionfish and <15 cm TL; Table S1, Supporting information).

In line with our predictions, vulnerability to predation, measured as the probability of being consumed, decreased significantly with prey size and with increasingly deep body shapes (Table 2; Fig. 1). Prey living solitarily or forming ephemeral feeding shoals were more likely to be consumed by lionfish than schooling prey, as were prey living on or in the benthos (Table 2; Fig. 1). We also found some evidence that fishes that exhibit cleaning behaviour were less vulnerable to predation than their non-cleaning counterparts (Table 2; Fig. 1). We observed no instances of predation on nocturnal, physically defended or pelagic prey; however, very few individuals of each of these types were observed within the vicinity of hunting lionfish on Eleuthera reefs (Table 2; Fig. 1).

PREY SELECTION FROM STOMACH CONTENTS ANALYSIS

In total 637 lionfish were collected in New Providence and 108 contained whole fish prey that could be identified to species, either visually or through DNA barcode analysis. From these specimens, we documented a total of 258 prey fishes from 33 species and 15 families (Table S1, Supporting information). The lionfish examined ranged in

sizes from 10 to 38 cm TL (25 ± 5 cm; mean \pm SD) and their fish prey ranged from 2 to 13 cm TL (4 ± 2 cm; mean \pm SD). The maximum number of prey fishes observed in a single lionfish stomach was 15; more than half (57 of 108) of the lionfish stomachs contained only one prey fish. We recorded 103 species from 30 families during the visual surveys of potential prey (i.e. all fishes <15 cm TL; Table S1, Supporting information).

On New Providence reefs, the vulnerability of prey fishes to lionfish predation again decreased with prey size and increasingly deep body shapes (Table 2; Figs 1 and 2). We found that cleaning behaviour predicted consumption by lionfish, with cleaners being selected relatively less frequently than non-cleaners (Table 2; Figs 1 and 2d). Aggregation behaviour and nocturnal activity also strongly influenced lionfish prey selection on New Providence reefs (Table 2; Figs 1 and 2b,c). As on Eleuthera reefs, schooling fishes were significantly less vulnerable than their solitary and shoaling counterparts, with solitary fishes most vulnerable to predation by lionfish (Table 2; Fig. 1). The presence of physical defence did not affect fish vulnerability to lionfish predation. Again, we observed no instances of predation on pelagic prey, despite their prevalence within our environmental surveys of New Providence reefs (Table 2; Fig. 1). When these traits are considered in combination, fishes that exhibit vulnerable forms of each morphological and behavioural trait are ~ 200 times more likely to be consumed by lionfish than their 'invulnerable' counterparts of the same size and body shape on New Providence reefs (Fig. 2).

Discussion


The majority of our predictions about the behavioural and morphological features of reef fishes that affect selection by lionfish predators were confirmed by data collected at two very different spatial scales, demonstrating that vulnerability to predation can be conferred by traits that recur across species within prey assemblages. In particular, our analyses of both *in situ* behavioural observations and *ex situ* stomach contents reveal that prey size, body shape, aggregation and cleaning behaviour are important determinants of vulnerability to predation, with small, solitary, and shallow-bodied fishes that do not engage in parasite removal consumed selectively. Water column position also strongly influenced vulnerability to predation; prey occurring on and in the benthos were most vulnerable to predation on Eleuthera reefs and pelagic prey occupying the water column >2 m above reefs were never selected by lionfish in our study. Nocturnality also predicted lionfish prey selection in our large-scale analysis of New Providence reefs.

Counter to our prediction, we found that schooling behaviour decreased the vulnerability of fish prey to lionfish predation on New Providence reefs and that solitary behaviour increased vulnerability on Eleuthera reefs,

Table 2. Top generalized linear mixed-effects models (GLMMs) of lionfish diet selection from (A) *in situ* observations of lionfish predation on coral reefs off Eleuthera, the Bahamas, and (B) inferred from lionfish stomach contents and visual surveys of reef fish on coral reefs off New Providence, the Bahamas. Δ AICc is the difference in AICc values between model *i* and the best model of those considered, and w_i is the probability that model *i* is the best of the set. The top models considered are those with Δ AICc < 4. Relative variable importance is the sum of the weights of all models that contain that particular variable

Response variable	Morphology				Behaviour				logLik	AICc	Δ AICc	Weight	df.
	Rank	Body shape	Body size	Cleaner	Defence	Aggregation size	Nocturnality	Water column position					
<i>In situ</i> feeding observations	1	✓	✓			✓			-69.41	157.90	0.00	0.19	9
	2	✓	✓			✓	✓		-69.09	159.50	1.60	0.09	10
	3	✓	✓		✓				-69.14	159.60	1.72	0.08	10
	4	✓	✓					✓	-69.24	159.90	1.92	0.07	10
	5	✓	✓		✓				-69.29	159.90	2.02	0.07	10
	6	✓	✓						-72.67	160.00	2.08	0.07	7
	7								-71.58	160.00	2.12	0.07	8
	8					✓			-70.73	160.60	2.64	0.05	9
	9					✓			-74.04	160.60	2.66	0.05	6
	10	✓				✓	✓		-68.80	161.30	3.32	0.04	11
	11					✓			-68.90	161.50	3.52	0.03	11
	12					✓			-68.92	161.50	3.56	0.03	11
	13				✓				-72.33	161.50	3.61	0.03	8
	14	✓			✓				-69.00	161.70	3.72	0.03	11
Stomach contents analysis	1	✓	✓	✓		✓			-370.78	762.00	0.00	0.32	10
	2	✓	✓	✓		✓			-369.91	762.30	0.33	0.27	11
	3	✓	✓	✓		✓			-374.08	764.40	2.44	0.10	8
	4	✓	✓	✓		✓			-375.27	764.80	2.76	0.08	7
	5	✓	✓	✓		✓			-373.25	764.90	2.86	0.08	9
	6	✓	✓	✓		✓			-372.45	765.30	3.33	0.06	10
	7	✓	✓	✓		✓			-374.74	765.80	3.76	0.05	8
	8	✓	✓	✓	✓	✓			-374.82	765.90	3.92	0.05	8
Relative variable importance (direct/indirect)		0.63/0.87	1/1	0.16/0.89	0.23/0.5	0.85/1	0.18/1	0.2/0.65					

Fig. 1. Model-averaged coefficients from the top generalized linear mixed-effects models (GLMMs; Δ AICc < 4; Table 2) of lionfish prey selection on Eleuthera coral reefs (a, *in situ* observations) and on New Providence coral reefs (b, stomach contents analysis). Points represent mean values bounded by 95% confidence intervals. Positive values (to the right of the vertical line) indicate selective predation on prey displaying the trait, while negative values (to the left of the vertical line) indicate avoidance. Levels 'demersal' and 'pelagic' are compared against the baseline level 'benthic', and 'schooling' and 'solitary' are compared against 'shoaling'. 'Body shape' is measured as the ratio of body length (cm) to depth (cm), and 'body size' is measured as mass (in g) as per Table 1. Numbers denoted with * are average parameter values, estimated with large uncertainty [i.e. standard error (SE) > 100] because no individuals with that specific trait form were consumed by lionfish in the data set.



supporting the hypothesis that schooling behaviour can be effective anti-predator strategy (Pitcher & Parrish 1993). While our observations indicate that on the whole, solitary species are more vulnerable than schooling species, the success of schooling behaviour as an anti-predator strategy has been shown to depend on a combination school size, prey vigilance, and predator detection strategy, which are thought to be species and context dependent (Bednekoff & Lima 1998). Further work is needed to tease apart the effects of school size and prey vigilance on vulnerability to lionfish predation. Another prey feature that unexpectedly had little influence on vulnerability to lionfish predation was the presence of a physical or chemical defence. One possible explanation is that the manner in which prey use physical defences (e.g. spines or barbs) may be ineffective against the hunting strategy of lionfish. During many of our direct observations, prey appeared largely unaware of stalking lionfish immediately prior to a capture. Alternatively, the defences of small-bodied prey may simply be ineffective against relatively large-bodied lionfish predators; prey were on average only 15% of the lionfish's length.

We show that mortality from lionfish predation will be greatest for prey possessing a specific suite of behavioural and morphological characteristics on invaded Atlantic coral reefs (e.g. Fig. 2). As lionfish can reach densities that allow the rapid depletion of prey biomass (Green *et al.* 2012), prey types that are selectively consumed may post more rapid and substantial declines over time, with potentially serious implications for local population persistence. Time series data documenting the relative change

in biomass of prey types over time, in relation to lionfish predation pressure, are needed to test this prediction. Moreover, selective predation by lionfish may have repercussions on invaded marine food webs if vulnerability to lionfish predation correlates with functional role of prey. For example, herbivorous fishes grow through size classes vulnerable to lionfish predation, and many species are of a body shape (body depth ratio range 4–7) that is vulnerable to predation by lionfish (Fig. 2a; Table S1, Supporting information). Thus, a high rate of lionfish-induced mortality on juvenile stages could impair the process of herbivory, which suppresses algal growth on coral reefs (Mumby *et al.* 2006), in the long term. Because our analyses focus on prey characteristics and not species identity, our findings may provide useful insight into the potential effects of lionfish predation on fish communities elsewhere in the invaded region; data on fish assemblages from before and early in the invasion of a region may be used to establish spatial priorities for management action. For example, locations where native fish communities are composed of a high proportion of vulnerable individuals (i.e. small-bodied, shallow-bodied, benthic or demersal, and nocturnal species) could be targeted for lionfish culling.

Predicting the strength of novel predator–prey is essential for understanding how ecological patterns and processes will be affected by the ongoing process of human-mediated species exchange. In particular, biological invasions are a major consequence of species exchange that are occurring at an increasing rate (Mooney & Cleland 2001). The expansion of the Burmese python (*Python*




Fig. 2. Example contour plots of predicted mean vulnerability to predation by lionfish for various combinations of prey size (mass in g) and body shape (quantified as the ratio of fish length to body depth; lower values indicate deeper-bodied shapes, whereas higher values denote shallow-bodied shapes) for fishes on coral reefs off New Providence, the Bahamas, displaying varying morphological and behavioural traits: (a) shoaling demersal fishes such as *Scarus iserti* and the deeper-bodied *Acanthurus bahianus*, (b) schooling pelagic fishes such as *Decapterus macarellus*, (c) solitary nocturnal fishes such as *Apogon binotatus* and (d) benthic cleaning fishes such as *Elacatinus genie*. Black contour lines denote a change in probability of predation. Cooler colours denote the combinations of prey fish size and body depth ratio yielding low vulnerability to predation, while warmer colours denote scenarios in which vulnerability to predation is high. Species illustrations modified with permission from ReefNet (2007).

molurus bivittatus) northwards from the Florida Everglades (Rodda, Jarnevic & Reed 2009), recent spread of Asian tiger prawn (*Penaeus monodon*) into the Gulf of Mexico and western Atlantic (USGS 2013) and progression of snakehead fishes (Channidae) throughout North American freshwater systems (Herborg *et al.* 2007) are just a few examples of non-native predators that are rapidly expanding into new areas, where they encounter and consume novel prey items. Low trophic level species are also invading non-native ranges at an increasing rate, where they alter the diet of native predators [e.g. the consumption of invasive gypsy moth (*Lymantria dispar*) by native cuckoos (*Coccyzus erythrophthalmus* and *Coccyzus americanus*) (Barber, Marquis & Tori 2008)]. Using our approach of identifying general prey traits that confer vulnerability to predation will aid in forecasting changes to predator–prey relationships, and ultimately consequences for community structure, in each of these systems. The only requirements to apply a trait-based approach are that (i) the predator's diet consists of multiple species, (ii) prey species share at least 1 trait that is predicted to affect vulnerability to predation and (iii) information on prey

consumption and availability be obtained, either through *in situ* observation of predation or by pairing broadscale diet information from guts or tissues with surveys of environmental prey abundance.

Finally, our study shows that a functional, or trait-based, approach has value for generating predictions about the effects of prey selection on the structure of ecological communities (i.e. McGill *et al.* 2006). Current thinking about predator–prey interactions can be broadly characterized into two prevailing paradigms: a species-based view, in which food webs are constructed by quantifying the interaction strength between pairs of predators and prey (Paine 1980; Polis & Strong 1996), and a size-based view, which classifies predator–prey interactions based on body size, largely ignoring species identity (Jennings 2005; Blanchard *et al.* 2011). Our study is the first to identify general traits of prey that predict vulnerability to predation and indicates that a trait-based view of predator–prey interactions, in which variation in morphology and behaviour determines prey selection, also has significant influence on species interactions within ecological communities (e.g. Hartvig, Andersen & Beyer 2011). It is

also, to our knowledge, the first to examine diet selection simultaneously at two spatial scales. Importantly, both our *in situ* observations of predation and broadscale analyses of stomach contents identified the same prey traits as important drivers of diet selection. However, directly observing prey capture *in situ* is challenging and information on diet composition is commonly gathered via methods similar to our broadscale analyses, with ‘snapshots’ of diet obtained by examining gut content (as we did) or scat composition. The similarity of traits identified by both analyses suggests that diet selection can be well estimated from these broadscale analyses when direct observation of predation is not logistically feasible.

Acknowledgements

We are grateful to Lad Akins, Lindsay Jacobs, Susan Reigner, Annabelle Brooks, Leah Neal and many Reef Environmental Education Foundation volunteers and Cape Eleuthera Institute interns for assistance in the field. We are also grateful to Mark Hixon, Ted Grosholz, Nicholas Dulvy, Jonathan Moore, John Bruno, Julia Baum and Julian Caley for comments on the manuscript and to Andrew Cooper and the Earth to Ocean Research Group for discussion of statistical analyses. Logistical support for this study on New Providence was generously donated by Stuart Cove’s Dive Bahamas and on Eleuthera by Trisha and David Ferguson. Funding was provided by a Natural Science and Engineering Research Council (NSERC) Canada Graduate Fellowship to SJG and an NSERC Discovery Grant to IMC.

Data accessibility

Data used in analyses: species data uploaded as online supporting information (Table S1, Supporting information) and available through FigShare pending acceptance of the study for publication.

References

- Albins, M.A. & Hixon, M.A. (2008) Invasive Indo-Pacific lionfish *Pterois volitans* reduce recruitment of Atlantic coral-reef fishes. *Marine Ecology Progress Series*, **367**, 233–238.
- Almany, G.R. & Webster, M.S. (2004) Odd species out as predators reduce diversity of coral-reef fishes. *Ecology*, **85**, 2933–2937.
- Almany, G., Peacock, L., Syms, C., McCormick, M. & Jones, G. (2007) Predators target rare prey in coral reef fish assemblages. *Oecologia*, **152**, 751–761.
- Arcese, P., Smith, J.N. & Hatch, M.I. (1996) Nest predation by cowbirds and its consequences for passerine demography. *Proceedings of the National Academy of Sciences*, **93**, 4608–4611.
- Barber, N.A., Marquis, R.J. & Tori, W.P. (2008) Invasive prey impacts the abundance and distribution of native predators. *Ecology*, **89**, 2678–2683.
- Barton, K. (2013) MuMIn: Multi-model inference. R package version 1.9.13. Available at: <http://CRAN.R-project.org/package=lme4>.
- Bates, D., Maechler, M. & Bolker, B. (2012) lme4: Linear Mixed-Effects Models Using Eigen and S4. R Package Version 0.999999-0. Available at: <http://CRAN.R-project.org/package=lme4>.
- Bednekoff, P.A. & Lima, S.L. (1998) Re-examining safety in numbers: interactions between risk dilution and collective detection depend upon predator targeting behaviour. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **265**, 2021–2026.
- Betancur, R.R., Hines, A., Acero, A.P., Ort, G.í., Wilbur, A.E. & Freshwater, D.W. (2011) Reconstructing the lionfish invasion: insights into Greater Caribbean biogeography. *Journal of Biogeography*, **38**, 1281–1293.
- Beukers-Stewart, B.D. & Jones, G.P. (2004) The influence of prey abundance on the feeding ecology of two piscivorous species of coral reef fish. *Journal of Experimental Marine Biology and Ecology*, **299**, 155–184.
- Blanchard, J., Law, R., Castle, M. & Jennings, S. (2011) Coupled energy pathways and the resilience of size-structured food webs. *Theoretical Ecology*, **4**, 289–300.
- Böhlke, J.E. & Chaplin, C.C. (1993) *Fishes of the Bahamas and Adjacent Tropical Waters*, 2nd edn. University of Texas Press, Austin, Texas.
- Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, H.H. *et al.* (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution*, **24**, 127–135.
- Bost, C.A., Zorn, T., Maho, Y.L. & Duhamel, G. (2002) Feeding of diving predators and diel vertical migration of prey: King penguins diet versus trawl sampling at Kerguelen Islands. *Marine Ecology Progress Series*, **227**, 51–61.
- Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: A Practical Information Theoretic Approach*. Springer, New York, New York.
- Caley, M.J. (1993) Predation, recruitment and the dynamics of communities of coral-reef fishes. *Marine Biology*, **117**, 33–43.
- Caley, M.J. & Schluter, D. (2003) Predation favors mimicry in a reef fish. *Proceedings of the Royal Society B: Biological Sciences*, **270**, 667–672.
- Carr, M.H. & Hixon, M.A. (1995) Predation effects on early postsettlement survivorship of coral-reef fishes. *Marine Ecology Progress Series*, **124**, 31–42.
- Caut, S., Angulo, E. & Courchamp, F. (2008) Dietary shift of an invasive predator: rats, seabirds and sea turtles. *Journal of Applied Ecology*, **45**, 428–437.
- Colin, P.L. (1975) *The Neon Gobies: The Comparative Biology of the Gobies of the Genus Gobiosoma, Subgenus Elacatinus*, (Pisces: Gobiidae) in the Tropical Western North Atlantic Ocean. T.F.H. Publications, Hong Kong.
- Côté, I.M. (2000) Evolution and ecology of cleaning symbioses in the sea. *Oceanography and Marine Biology*, **38**, 311–355.
- Côté, I.M. & Maljković, A. (2010) Predation rates of Indo-Pacific lionfish on Bahamian coral reefs. *Marine Ecology Progress Series*, **404**, 219–225.
- Côté, I.M., Green, S.J. & Hixon, M.A. (2013b) Predatory fish invaders: insights from Indo-Pacific lionfish in the western Atlantic and Caribbean. *Biological Conservation*, **164**, 50–61.
- Côté, I.M., Green, S.J., Morris, J.A.J., Akins, J.L. & Steinke, D. (2013a) Diet richness of an invasive Indo-Pacific lionfish revealed by DNA barcoding. *Marine Ecology Progress Series*, **472**, 249–256.
- Creel, S., Christianson, D., Liley, S. & Winnie, J.A. (2007) Predation risk affects reproductive physiology and demography of elk. *Science*, **315**, 960.
- Crooks, K. & Soulé, M. (1999) Mesopredator release and avifaunal extinctions in a fragmented system. *Nature*, **400**, 563–566.
- Cunningham, S.C., Gustavson, C.R. & Ballard, W.B. (1999) Diet selection of mountain lionfish in southeastern Arizona. *Journal of Range Management*, **52**, 202–207.
- Deloach, N. (1999) *Reef Fish Behavior: Florida, Caribbean and Bahamas*. New World Publications Inc., Jacksonville, Florida, USA.
- Donazar, J. & Ceballos, O. (1989) Selective predation by eagle owls *Bubo bubo* on rabbits *Oryctolagus cuniculus*: age and sex preferences. *Ornis Scandinavica*, **20**, 117–122.
- Einfalt, L.M. & Wahl, D.H. (1997) *Prey Selection by Juvenile Walleye as Influenced by Prey Morphology and Behavior*. National Research Council of Canada, Ottawa, ON, Canada.
- FAO. (2002a) Bony fishes part 1 (Acipenseridae to Grammatidae). *The Living Marine Resources of the Western Central Atlantic* (eds K.E. Carpenter), pp. 601–1374. FAO, Rome.
- FAO. (2002b) Bony fishes part 2 (Opistognathidae to Molidae), sea turtles and marine mammals. *The Living Marine Resources of the Western Central Atlantic* (ed. K.E. Carpenter), pp. 1375–2127. FAO, Rome.
- Fishelson, L. (1975) Ethology and reproduction of pteroid fishes found in the Gulf of Aqaba (Red Sea), especially *Dendrochirus brachypterus* (Cuvier), (Pteroidae, Teleostei). *Pubblicazioni della Stazione Zoologica di Napoli*, **39**, 635–656.
- Gelman, A. (2008) Scaling regression inputs by dividing by two standard deviations. *Statistics in Medicine*, **27**, 2865–2873.
- Gilinsky, E. (1984) The role of fish predation and spatial heterogeneity in determining benthic community structure. *Ecology*, **65**, 455–468.
- Green, S.J., Akins, J.L. & Côté, I.M. (2011) Foraging behaviour and prey consumption in the Indo-Pacific lionfish on Bahamian coral reefs. *Marine Ecology Progress Series*, **433**, 159–167.
- Green, K. & Osborne, W. (1981) The diet of foxes, *Vulpes vulpes* (L.), in relation to abundance of prey above the winter snowline in New South Wales. *Wildlife Research*, **8**, 349–360.

- Green, S.J., Akins, J.L., Maljković, A. & Côté, I.M. (2012) Invasive lionfish drive Atlantic coral reef fish declines. *PLoS One*, **7**, e32596.
- Green, S.J., Dulvy, N.K., Brooks, A., Akins, J.L., Cooper, A.B., Miller, S. *et al.* (in press) Linking removal targets to the ecological effects of invaders: a predictive model and field test. *Ecological Applications*, doi:10.1890/13-0979.1.
- Grueber, C.E., Nakagawa, S., Laws, R.J. & Jamieson, I.G. (2011) Multi-model inference in ecology and evolution: challenges and solutions. *Journal of Evolutionary Biology*, **24**, 699–711.
- Hambright, K.D. (1991) Experimental-analysis of prey selection by large-mouth bass- role of predator mouth width and prey body depth. *Transactions of the American Fisheries Society*, **120**, 500–508.
- Hartvig, M., Andersen, K.H. & Beyer, J.E. (2011) Food web framework for size-structured populations. *Journal of Theoretical Biology*, **272**, 113–122.
- Heinonen, K.B. & Auster, P.J. (2012) Prey selection in crustacean-eating fishes following the invasion of the Asian shore crab *Hemigrapsus sanguineus* in a marine temperate community. *Journal of Experimental Marine Biology and Ecology*, **413**, 177–183.
- Herborg, L.-M., Mandrak, N.E., Cudmore, B.C. & MacIsaac, H.J. (2007) Comparative distribution and invasion risk of snakehead (Channidae) and Asian carp (Cyprinidae) species in North America. *Canadian Journal of Fisheries and Aquatic Sciences*, **64**, 1723–1735.
- Hernández, A. (1995) Selective predation by Northern Shrikes on small mammals in a natural environment (Depredación Selectiva Sobre Micromamíferos por Lanius excubitor en Condiciones Naturales). *Journal of Field Ornithology*, **66**, 236–246.
- Holt, R.D. (1984) Spatial heterogeneity, indirect interactions, and the coexistence of prey species. *The American Naturalist*, **124**, 377–406.
- Hoogland, R., Morris, D. & Tinbergen, N. (1956) The spines of sticklebacks (*Gasterosteus* and *Pygosteus*) as means of defence against predators (*Perca* and *Esox*). *Behaviour*, **10**, 205–236.
- Humann, P. & Deloach, N. (2002) *Reef Fish Identification: Florida, Caribbean and Bahamas*, 3rd edn. New World Publications Inc., Jacksonville, Florida, USA.
- Isaac, E.J., Hrabik, T.R., Stockwell, J.D. & Gamble, A.E. (2012) Prey selection by the Lake Superior fish community. *Journal of Great Lake Research*, **38**, 326–335.
- Jaeger, A., Connan, N., Richard, P. & Cherel, Y. (2010) Use of stable isotopes to quantify seasonal changes of trophic niche and levels of population and individual specialisation in seabirds. *Marine Ecology Progress Series*, **401**, 269–277.
- Jennings, S., Greenstreet, S.P.R. & Reynolds, J.D. (1999) Structural change in an exploited fish community: a consequence of differential fishing effects on species with contrasting life histories. *Journal of Animal Ecology*, **68**, 617–627.
- Jennings, S. (2005) Size-based analyses of aquatic food webs. *Aquatic Food Webs: An Ecosystem Approach* (eds A. Belgrano, U. Scharler, J. Dunne & R. Ulanowicz), pp. 86–97. Oxford University Press, Oxford.
- Juanes, F., Buckel, J.A. & Scharf, F.S. (2001) Predatory behaviour and selectivity of a primary piscivore: comparison of fish and non-fish prey. *Marine Ecology Progress Series*, **217**, 157–165.
- Karanth, K.U. & Sunquist, M.E. (1995) Prey selection by tiger, leopard and dhole in tropical forests. *Journal of Animal Ecology*, **64**, 439–450.
- Kotler, B.P. (1984) Risk of predation and the structure of desert rodent communities. *Ecology*, **65**, 689–701.
- Lima, M., Julliard, R., Stenseth, N.C.H.R. & Jaksic, F.M. (2001) Demographic dynamics of a neotropical small rodent (*Phyllotis darwini*): feedback structure, predation and climatic factors. *Journal of Animal Ecology*, **70**, 761–775.
- McGill, B.J., Enquist, B.J., Weiher, E. & Westoby, M. (2006) Rebuilding community ecology from functional traits. *Trends in Ecology and Evolution*, **21**, 178–185.
- Mooney, H.A. & Cleland, E.E. (2001) The evolutionary impact of invasive species. *Proceedings of the National Academy of Sciences of the United States of America*, **98**, 5446–5451.
- Morris, J.A.J. & Akins, J.L. (2009) Feeding ecology of invasive lionfish (*Pterois volitans*) in the Bahamian Archipelago. *Environmental Biology of Fishes*, **86**, 389–398.
- Mumby, P.J., Dahlgren, C.P., Harborne, A.R., Kappel, C.V., Micheli, F., Brumbaugh, D.R. *et al.* (2006) Fishing, Trophic Cascades, and the Process of Grazing on Coral Reefs. *Science*, **311**, 98–101.
- Munoz, R., Currin, C. & Whitfield, P. (2011) Diet of invasive lionfish on hard bottom reefs of the Southeast USA: insights from stomach contents and stable isotopes. *Marine Ecology Progress Series*, **432**, 181–193.
- Myers, R.A. & Worm, B. (2003) Rapid worldwide depletion of predatory fish communities. *Nature*, **423**, 280–283.
- Nilsson, P.A. & Brönmark, C. (2000) Prey vulnerability to a gape-size limited predator: behavioural and morphological impacts on northern pike piscivory. *Oikos*, **88**, 539–546.
- Nunn, A.D., Tweson, L.H. & Cowx, I.G. (2012) The foraging ecology of larval and juvenile fishes. *Reviews in Fish Biology and Fisheries*, **22**, 377–408.
- Paine, R.T. (1980) Food Webs: linkage, interaction strength and community infrastructure. *Journal of Animal Ecology*, **49**, 667–685.
- Phillips, R., Bearhop, S., McGill, R.R. & Dawson, D. (2009) Stable isotopes reveal individual variation in migration strategies and habitat preferences in a suite of seabirds during the nonbreeding period. *Oecologia*, **160**, 795–806.
- Pitcher, T. & Parrish J. K. (1993) Antipredator functions of shoaling. *The Behaviour of Teleost Fishes* (ed. T. Pitcher), pp. 379–400. Croom Helm Ltd., London.
- Polis, G.A. & Strong, D.R. (1996) Food web complexity and community dynamics. *American Naturalist*, **147**, 813–846.
- Provost, C., Lucas, E., Coderre, D. & Chouinard, G. (2006) Prey selection by the lady beetle *Harmonia axyridis*: the influence of prey mobility and prey species. *Journal of Insect Behavior*, **19**, 265–277.
- Pyke, G.H., Pulliam, H.R. & Charnov, E.L. (1977) Optimal foraging: a selective review of theory and tests. *The Quarterly Review of Biology*, **52**, 137.
- R Core Team (2012) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- ReefNet (2007) *Reef Fish Identification: Florida, Caribbean, Bahamas*. ReefNet Inc., Ontario, Canada.
- Rodda, G., Jarnevech, C. & Reed, R. (2009) What parts of the US mainland are climatically suitable for invasive alien pythons spreading from Everglades National Park? *Biological Invasions*, **11**, 241–252.
- Scharf, F.S., Buckel, J.A., McGinn, P.A. & Juanes, F. (2003) Vulnerability of marine forage fishes to piscivory: effects of prey behavior on susceptibility to attack and capture. *Journal of Experimental Marine Biology and Ecology*, **294**, 41–59.
- Schubert, M.J., Munday, P.L., Caley, M.J., Jones, G.P. & Llewellyn, L.E. (2003) The toxicity of skin secretions from coral-dwelling gobies and their potential role as a predator deterrent. *Environmental Biology of Fishes*, **67**, 359–367.
- Scott, M.A. & Murdoch, W.W. (1983) Selective predation by the backswimmer, *Notonecta*. *Limnology and Oceanography*, **28**, 352–366.
- Steele, J. (1989) The ocean 'landscape'. *Landscape Ecology*, **3**, 185–192.
- USGS (2013) Non-indigenous Aquatic Species Database. U.S. Geological Survey. <http://nas.er.usgs.gov/> Accessed online December 2013.
- Wen, C.K., Almany, G.R., Williamson, D.H., Pratchett, M.S. & Jones, G.P. (2012) Evaluating the effects of marine reserves on diet, prey availability and prey selection by juvenile predatory fishes. *Marine Ecology Progress Series*, **469**, 133–144.
- Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A. & Smith, G.M. (2009) *Mixed Effects Models and Extensions in Ecology with R*. Springer, New York, USA.

Received 6 February 2014; accepted 8 May 2014

Handling Editor: Stan Boutin

Supporting Information

Additional Supporting Information may be found in the online version of this article.

Table S1. List of fish species observed on Bahamian coral reefs and their morphological and behavioural characteristics which are hypothesized to affect vulnerability to predation by invasive lionfish.